

(12) UK Patent Application (19) GB (11) 2 031 410 A

- (21) Application No 7929266
(22) Date of filing 22 Aug 1979
(23) Claims filed 22 Aug 1979
(30) Priority data
(31) 7824674
(32) 25 Aug 1978
(33) France (FR)
(43) Application published
23 Apr 1980
(51) INT CL³

C07D 265/26 A61K
31/535 C07D 413/02
(C07D 413/02 213/72
265/26 307/52)
(52) Domestic classification
C2C 1470 1564 213 215
220 226 22Y 246 247 255
26Y 28X 305 313 314
31Y 337 351 352 364 38Y
373 37Y 386 388 401 40Y
553 613 624 635 662 672
699 781 768 775 802 80Y
AA OT TX

- (56) Documents cited
GB 1195232
GB 1131467
GB 971246
GB 950065
GB 817147
US 3443014A
US 2476569A
Eur J Med. Chem-Chim
Ther 9 44-50 (1974) J Het
Chem 2 37-40 (1965)

- (58) Field of search
C2C


- (71) Applicants
Provesan S.A., 1, Place
St. Gervais, 1211
Geneva, Switzerland

- (72) Inventor
Antonio Esteve-Subirana

- (74) Agents
Reddie & Grose

(54) New derivatives of 1,3-benzoxazine-2,4-dione, their preparation and their application as medicaments

(57) The invention concerns 1,3-benzoxazine-2, 4-dione derivatives of the formula



In which R represents an alkyl radical, an *ortho*-phenoxy-phenyl, *ortho*-thiophenoxy-phenyl, a benzyl, furfuryl or 2-pyridyl group; R' represents a hydrogen, chlorine, or bromine atom or a methoxy radical; R'' represents a hydrogen or chlorine atom or a methyl radical; and R''' represents a hydrogen, chlorine or bromine atom. R preferably represents a C₁ to C₄ alkyl radical. The derivatives of the invention have pharmacological (e.g. analgesic and anti-inflammatory) activity and may be administered in tablet, capsule, suppository and the like forms. Methods for the preparation of these derivatives are disclosed.

Certain of the chemical formulae appearing in the printed specification were submitted in formal form after the date of filing.

BEST AVAILABLE COPY


GB 2 031 410 A

SPECIFICATION

New derivatives of 1,3-benzoxazine-2,4-dione, their preparation and their application as medicaments

The present invention provides new derivatives of 1,3-benzoxazine-2,4-dione, their preparation and their application as medicaments.

The new derivatives of the present invention, have with the general formula I



in which R represents an alkyl radical, especially a lower alkyl radical preferably with C₁ to C₄, an ortho-phenoxy-phenyl, ortho-thiophenoxy-phenyl, a benzyl, furfuryl or a 2-pyridyl group;

R' represents a hydrogen, chlorine, or bromine atom or a methoxyl radical;

R'' represents a hydrogen or chlorine atom or a methyl radical; and


R''' represents a hydrogen, chlorine or bromine atom.

The derivatives of general formula I have proved to possess valuable pharmacological properties.

The present invention therefore also relates to the application of these compounds as medicaments as well as to pharmaceutical compositions containing them as active ingredient.


The present invention also relates to the preparation of derivatives of general formula I. According to the invention, these derivatives of general formula I are obtained:

a) by reaction of a derivative of salicylamide of general formula II



In which R, R', R'' and R''' have the meanings indicated in connection with general formula I, with ethyl chlorocarbonate; or

b) by reaction of a derivative of salicylic acid of general formula III



in which R', R'' and R''' have the meaning indicated in connection with general formula I, with an isocyanate of general formula IV



in which R has the meaning given in connection with general formula I.

By way of simple non-limiting examples, the preparation of a few derivatives of general formula I will hereinafter be described in more detail.

EXAMPLE 1***Preparation of 6-chloro-3-methyl-1, 3-benzoxazine-2, 4-dione***

37 ml (0.375 mole) of ethyl chlorocarbonate is added slowly (1/2 hour) to a solution of 18.6 g (0.1 mole) of 5-chloro-N-methylsalicylamide in pyridine. It is allowed to reflux for 7 hours, allowed to cool, is diluted with water, filtered and washed with distilled water. It is recrystallised from chloroform, washed with methanol at 5°C and 6-chloro-3-methyl-1, 3-benzoxazine-2, 4-dione with melting point 151—153°C (see Table I) is obtained.

EXAMPLE 2***Preparation of 6-bromo-3-methyl-1, 3-benzoxazine-2, 4-dione***


7.5 ml (0.12 mole) of methyl isocyanate and 5 ml of triethylamine are added to a solution of 21.7 g (0.1 mole) of 5-bromosalicylic acid in 150 ml of benzene.

The mixture is stirred for 1 hour at ambient temperature and then allowed to reflux for 6 hours. It is evaporated to dryness, recrystallised from methanol and 6-bromo-3-methyl-1,3-benzoxazine-2,4-dione with melting point 186°C (see Table I) is obtained.

EXAMPLES 3 TO 17


By operating as described in Examples 1 and 2 various derivatives were obtained, a number of the physico-chemical constants of which, such as the crystallisation solvents, the melting points and the characteristic bands of the infra-red spectrum are indicated in Table I hereinafter.

TABLE I



Ex. No.	R	R'	R''	R'''	Melting Point	Crystallization Solvent	Yield %	IR (cm ⁻¹)	ANALYSIS					
									C	H	Cl	Br	N	
1	CH ₃	Cl	H	H	161-3	CHCl ₃	93	1300, 1360 1680, 1770	Calc. 51.12 Found 51.20	2.88 2.79	16.76 16.70	-	8.63 8.57	
2	CH ₃	Br	H	H	186	CH ₃ OH	19	1300, 1360 1690, 1750	Calc. 42.23 Found 42.15	2.36 2.41	-	31.14 31.27	5.47 5.32	
3	CH ₂ CH ₂ CH ₃	Cl	H	H	118	CH ₃ OH	52	1340 1700, 1770	Calc. 55.17 Found 55.25	4.21 4.15	14.80 14.73	-	5.85 5.97	
4	CH(CH ₃) ₂	Cl	H	H	167-9	CH ₃ OH	24	1330 1700, 1770	Calc. 55.17 Found 55.08	4.21 4.32	14.80 14.86	-	5.85 5.80	
5	CH ₂ CH ₂ CH ₂ CH ₃	Cl	H	H	103	CH ₃ CH ₂ OH	12	1300 1700, 1770	Calc. 56.87 Found 56.79	4.75 4.81	14.00 13.92	-	5.52 5.46	
6	(CH ₂) ₁₈ CH ₃	Cl	H	H	97	CH ₃ COCH ₃	67	1350 1700, 1770	Calc. 68.31 Found 68.42	8.60 8.56	8.40 8.35	-	3.32 3.25	
7		Cl	H	H	152-5	CH ₃ OH	42	1340 1710, 1770	Calc. 62.60 Found 62.55	3.51 3.57	12.33 12.28	-	4.87 4.97	
8		Cl	H	H	178-80	EtOH/Me ₂ CO	46	1320 1700, 1770	Calc. 56.27 Found 56.14	2.90 2.87	12.77 12.69	-	5.05 4.98	
9		Cl	H	H	167	EtOH	70	1360 1710, 1770	Calc. 65.73 Found 65.61	3.30 3.34	9.70 9.77	-	3.63 3.95	

TABLE I (continued)



Ex. No.	R	R'	R''	R'''	Melting Point	Crystallisation Solvent	Yield %	IR (cm ⁻¹)	ANALYSIS					
									C	H	Cl	Br	N	
10		Cl	H	H	150	EtOH	98	1380 1710, 1770	Calc. 62,90 Found 63,02	3,17 3,09	9,29 9,34	-	3,67 3,80	
11	CH ₃	H	Cl	H	182-4	CH ₃ OH	10	1300, 1360 1690, 1780	Calc. 51,11 Found 50,98	2,86 2,93	16,78 16,89	-	6,62 6,63	
12	CH ₃ CH ₃	Cl	H	H	120-2	C Cl ₄	51	1340, 1360 1700, 1770	Calc. 53,27 Found 53,15	3,58 3,63	16,72 15,74	-	6,22 6,15	
13	CH ₃	Cl	H	Cl	159-63	AcOEt	6	1300, 1360 1690, 1770	Calc. 43,84 Found 44,05	2,05 1,98	28,82 28,91	-	5,69 5,75	
14	CH ₃ CH ₂ CH ₂ CH ₃	Cl	H	Cl	106-8	CH ₃ OH	46	1340 1700, 1770	Calc. 50,00 Found 49,87	3,85 3,91	24,61 24,57	-	4,88 4,79	
15	CH ₃ CH ₂ CH ₂ CH ₂ CH ₃	Br	H	Br	139	EtOH	38	1350 1690, 1770	Calc. 38,23 Found 36,32	2,94 3,01	-	42,39 42,53	3,72 3,77	
16	CH ₃	CH ₃ O	H	H	138-40	CH ₃ OH/H ₂ O	23	1300, 1355 1690, 1780	Calc. 55,02 Found 57,91	4,38 4,42	-	-	6,77 6,85	
17	CH ₃	H	CH ₃	H	164	CH ₃ OH	12	1300, 1370 1690, 1760	Calc. 62,89 Found 62,97	4,75 4,70	-	-	7,33 7,40	

Analgesic activity

The analgesic activity of the derivatives of general formula I has been determined using male mice of weight between 20 and 25 g. The product to be tested is administered orally in suspension in 5 % gum arabic by means of an oesophageal probe. The volume of the solution administered is 25 ml/kg and the concentration of the tested product is changed according to the dose administered. 5

Pain is caused in the animals by an intraperitoneal injection of 0.2 ml/20g solution of acetylcholine bromide with a concentration of 0.32 ml/ml. Five minutes before the administration of the tested product, the acetylcholine is injected into a batch of 5 mice. The product to be tested is then administered and the injection of acetylcholine is administered again after 20, 40, 80, 120 and 160 minutes. The number of contortions per injection of acetylcholine is always counted for 5 minutes. 10

The analgesic activity is calculated by means of the following formula:

$$It = 100 - (Nt/No) \cdot 100 = 100 (1 - Nt/No)$$

where

It = inhibition of the pain after t minutes
15 No = number of contortions before the administration of the product 15

Nt = number of contortions after t minutes from the administration of the product.

Several doses of each product are administered so that the fifty per cent analgesic dose (AD—50) can be determined.

With each of these doses It is calculated for times of 20, 40, 80, 120 and 160 minutes. The mean 20 of these five values of It for each dose is taken as the analgesic effect. The analgesic effects are 20 represented graphically as a function of the logarithm of the corresponding dose.

From this curve the analgesic dose fifty, that is to say, the dose which produces a fifty per cent analgesic effect, is obtained.

By way of example, the results obtained for a few derivatives of formula I according to the invention 25 have been indicated in Table II hereinafter. 25

Acute toxicity

The acute toxicity is determined orally with mice of 20 to 25 g weight, by using batches of 6 animals. Several doses in geometric progression are administered. The time of observation is 72 hours. The fifty per cent lethal dose (LD—50) is calculated graphically by means of logarithmic-probabilistic 30 paper. 30

By way of example, the results obtained for a few derivatives of formula I according to the invention are indicated in Table II hereinafter.

TABLE II

Example No.	Derivative:	Doses in mg/kg	
		AD—50	LD—50
1	6-chloro-3-methyl-1,3-benzoxazine-2,4-dione	30	233
2	6-bromo-3-methyl-1,3-benzoxazine-2,4-dione	90	717
3	6-chloro-3-propyl-1,3-benzoxazine-2,4-dione	240	>650
8	6-chloro-3-furfuryl-1,3-benzoxazine-2,4-dione	280	>550
9	6-chloro-3-(O-phenoxyphenyl)-1,3-benzoxazine-2,4-dione	675	>750
10	6-chloro-3-(O-thiophenoxy-phenyl)-1,3-benzoxazine-2,4-dione	650	>1500
12	6-chloro-3-ethyl-1,3-benzoxazine-2,4-dione	172	>250

Anti-inflammatory activity

The anti-inflammatory activity in the male rat of Sprague-Dawley stock is determined. An oedema is caused in the paw by subplantar injection of a 1 % solution of carragheenin. The volume of the paw is measured before the oral administration of the product after two and five hours with a plethysmometer.

- 5 The anti-inflammatory activity is calculated with respect to a reference batch. By way of example, the results obtained for the derivative of example I are indicated in Table III. 5

TABLE III

Example	Derivative	Dose (mg/kg)	Anti-Inflammatory Activity	
			2 hours	5 hours
1	6-chloro-3-methyl- 1,3-benzoxazine- 2,4-dione	100	26%	30%

- Taking into account their good pharmacodynamic properties, the derivatives of general formula I are hence capable of being used as veterinary and/or human medicine, as analgesic, antipyretic and
10 anti-inflammatory agents. 10

- Pharmaceutical compositions which contain, according to the invention, besides an acceptable pharmaceutical support, at least one derivative of general formula I have a very large field of therapeutic application and can be utilised especially in traumatology, surgery, rheumatology, odontostomatology, oto-rhino-laryngology, pneumology, cardiology, gynaecology and urology. These pharmaceutical
15 compositions will be, for example, utilised for the treatment of various manifestations of pain, 15
headaches, migraines, toothache, neuralgias, menstrual pains, inflammatory rheumatisms, arthritis
pains, feverish states, colds, Influenzas and seasonal infections.

- In human therapy, the dose proposed for the derivatives of the present invention is approximately
20 between 100 and 300 mg/day, administered for example in the form of tablets, capsules or 20
suppositories.

Hereinafter, by way of example, three particular galenic forms of the derivatives, the objects of the present invention, will be indicated.

Example of formula per tablet

- | | | | |
|----|---|--------|----|
| 25 | 6-chloro-3-methyl-1, 3-benzoxazine-2, 4-dione | 100 mg | |
| | Avicel pH—102 | 100 mg | 25 |
| | Primojel | 10 mg | |
| | Aerosil—200 | 1 mg | |
| | Magnesium stearate | 2 mg | |

Tablet weight 213 mg

- 30 **Example of formula per capsule** 30
- | | | | |
|----|---|--------|----|
| | 6-chloro-3-methyl-1, 3-benzoxazine-2, 4-dione | 100 mg | |
| | Lactose | 75 mg | |
| | Avicel pH—102 | 25 mg | |
| | Aerosil—200 | 1 mg | |
| 35 | Magnesium stearate | 2 mg | 35 |


Capsule weight 203 mg

Example of formula per suppository

- | | | | |
|----|---|-------|----|
| | 6-chloro-3-methyl-1, 3-benzoxazine-2, 4-dione | 0.2 g | |
| | Monolene | 1.8 g | |
| 40 | Suppository weight | 2.0 g | 40 |

CLAIMS

1. A derivative of 1,3-benzoxazine-2,4-dione of the general formula



(I)

In which R represents an alkyl radical, an ortho-phenoxy-phenyl, ortho-thiophenoxy-phenyl, a benzyl, furfuryl or 2-pyridyl group; R' represents a hydrogen, chlorine, or bromine atom or a methoxy radical; R'' represents a hydrogen or chlorine atom or a methyl radical; and R''' represents a hydrogen, chlorine or bromine atom.

2. A derivative as claimed in claim 1 wherein R represents a lower alkyl radical.

3. A derivative as claimed in claim 1 wherein R represents an alkyl radical with 1 to 4 carbon atoms.

4. 6-chloro-3-methyl-1, 3-benzoxazine-2, 4-dione.

5. 6-chloro-3-ethyl-1, 3-benzoxazine-2, 4-dione.

6. 6-chloro-3-propyl-1, 3-benzoxazine-2, 4-dione.


7. 6-chloro-3-furfuryl-1, 3-benzoxazine-2, 4-dione.

8. 6-chloro-3-(o-phenoxyphenyl)-1, 3-benzoxazine-2, 4-dione.

9. 6-chloro-3-(o-thiophenoxyphenyl)-1, 3-benzoxazine-2, 4-dione.


10. 6-bromo-3-methyl-1, 3-benzoxazine-2, 4-dione.

11. A method of preparation of a derivative as claimed in claim 1, wherein a derivative of salicylamide of the general formula




in which R, R', R'' and R''' have the meaning indicated in claim 1 is reacted with ethyl chlorocarbonate.

12. A method of preparation of a derivative as claimed in claim 1, wherein a derivative of salicylic acid of the general formula



25 in which R', R'' and R''' have the meaning indicated in claim 1, is reacted with an isocyanate of general formula



in which R has the meaning given in claim 1.

13. A method as claimed in claim 14, conducted substantially as described in Example 1.

14. A method as claimed in claim 15, conducted substantially as described in Example 2.

15. A pharmaceutical composition comprising a compound as claimed in any one of claims 1 to 13 together with a pharmaceutically acceptable carrier.

16. A pharmaceutical composition as claimed in claim 19 in the form of a tablet, capsule or suppository.

5 17. A pharmaceutical composition as claimed in claim 19 substantially as described in the Examples herein.

5

Printed for Her Majesty's Stationery Office by the Courier Press, Leamington Spa, 1980. Published by the Patent Office,
25 Southampton Buildings, London, WC2A 1AY, from which copies may be obtained.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.